

FAMILIES OF SUPERSINGULAR CURVES IN CHARACTERISTIC 2

JASPER SCHOLTEN AND HUI JUNE ZHU

ABSTRACT. This paper determines all normal form of hyperelliptic supersingular curves of genus g over an algebraically closed field F of characteristic 2 for $1 \leq g \leq 8$. We also show that every hyperelliptic supersingular curve of genus 9 over F has an equation $y^2 - y = x^{19} + c^8 x^9 + c^3 x$ for some $c \in \overline{\mathbb{F}}_2$. Consequently, the paper determines the dimensions of the open locus of hyperelliptic supersingular curves of genus $g \leq 9$ over $\overline{\mathbb{F}}_2$.

1. INTRODUCTION

In this paper, a curve is a projective, smooth and geometrically integral algebraic variety of dimension 1. A curve is *supersingular* if its Jacobian is a supersingular abelian variety, that is, its Newton polygon is a straight line segment of slope $1/2$. See Introduction and Appendix of [6] for literatures on supersingular abelian varieties and related open questions on stratification in the moduli space of abelian varieties. It has been proved that there are supersingular curves of every genus over $\overline{\mathbb{F}}_2$ (see [10]), but it is not settled if there are hyperelliptic supersingular curves over $\overline{\mathbb{F}}_2$ of every genus, which was an initial goal of our study (see [7]). Several isolated discoveries of supersingular curves have yielded unprecedented applications in sphere packing [2] and [3]. There are also rich literatures in coding theorem upon applications of supersingular curves (see [11]). See the introduction in [7] for more recent progress regarding the hyperelliptic supersingular curves in characteristic 2. In [4] a question is raised that for which p and g there are positive dimensional families of hyperelliptic supersingular curves over $\overline{\mathbb{F}}_p$ of genus g . We give an answer to this question for $p = 2$ and $g \leq 9$ in Theorem 1.

Every genus- g hyperelliptic curve X over F of 2-rank zero has an affine equation

$$(1) \quad X : y^2 - y = c_{2g+1}x^{2g+1} + c_{2g-1}x^{2g-1} + \dots + c_1x,$$

where $c_1, \dots, c_{2g-1} \in F$ and $c_{2g+1} = 1$. (See Proposition 10).

For any g , an equation (1) with some vanishing conditions among c_ℓ 's is called a *normal form of hyperelliptic supersingular curve of genus g* if every such equation defines a hyperelliptic supersingular curve and conversely every hyperelliptic supersingular curve of genus g has such an equation for some c_ℓ 's in F , and there are only finitely many such equations for each curve.

Let $\mathcal{HS}_g/F$ denote the open locus of hyperelliptic supersingular curves of genus g over F .

Date: February 1, 2008.

1991 Mathematics Subject Classification. 11L, 14H, 14M.

Key words and phrases. supersingular curves, hyperelliptic curves, normal forms, moduli space of curves.

Theorem 1. *Let F be an algebraically closed field of characteristic 2. The normal forms of hyperelliptic supersingular curves over F of genus $g \leq 8$ are in the table below with all $c_\ell \in F$. The dimensions of $\mathcal{HS}_g/F$ are in the right columns.*

genus g	normal form X_g	$\dim(\mathcal{HS}_g/F)$
1	$y^2 - y = x^3$	0
2	$y^2 - y = x^5 + c_3x^3$	1
3	none	$-\infty$
4	$y^2 - y = x^9 + c_5x^5 + c_3x^3$	2
5	$y^2 - y = x^{11} + c_3x^3 + c_1x$	2
6	$y^2 - y = x^{13} + c_3x^3 + c_1x$	2
7	none	$-\infty$
8	$y^2 - y = x^{17} + c_9x^9 + c_5x^5 + c_3x^3$	3

TABLE 1. normal forms for $g \leq 8$

Moreover, every supersingular curve over F of genus 9 has an equation

$$y^2 - y = x^{19} + c^8x^9 + c^3x$$

for some $c \in \overline{\mathbb{F}}_2$. We have $\dim(\mathcal{HS}_9/F) = 0$.

Remark 2. From Theorem 1 one can easily derive the following: If $g = 1$ or 2 , then all 2-rank zero curves are supersingular. If $g = 3$ or 7 , then none of these curves is supersingular. If $g = 4$ then X is supersingular if and only if $c_7 = 0$. If $g = 5$ then X is supersingular if and only if $c_7 = 0$ and $c_9c_{11} = c_5^4$. If $g = 6$ then X is supersingular if and only if $c_7 = 0$, $c_{11} = 0$, and $c_5c_{13} = c_9^2$. If $g = 8$ then X is supersingular if and only if $c_{15} = c_{13} = c_{11} = c_7 = 0$.

Remark 3. It is clear from Theorem 1 that $\mathcal{HS}_g/F$ for $1 \leq g \leq 8$ are irreducible and unirational. One wonders if it is so for all g .

The proof of Theorem 1 splits into two parts: In section 4 we show that all curves in Table 1 are supersingular. In section 5 we show that every supersingular curve of genus ≤ 9 has an equation in Table 1.

One main idea in the proof of Theorem 1 is to use a lemma of [8] to reduce the initial problem to a combinatorial puzzle. Then we formulate an algorithm, utilizing Proposition 10 and Theorem 11 from [8], to prove our assertion.

We anticipate some results of similar sort for Artin-Schreier curves in characteristic p if p is small compared to the genus. For some results for p large, see [12].

Acknowledgments. We are grateful to Noam Elkies for his suggestion in section 4. We especially thank Bjorn Poonen for his valuable suggestions in Remark 2. The computation in this paper used Magma and Maple packages and C-programs.

2. 2-ADIC BOX AND A TILING PUZZLE

Let $s(\cdot)$ denote the number of 1's in the binary expansion of a positive integer. This section digresses to some new combinatorial notion. Let S be a finite set of positive integers. Let r be a positive integer. For a finite sequence of integer pairs $\{[b_i, \ell_i]\}_{i=1}^r$ such that

Conversely, given $\mathbf{k} \in \mathbf{K}_r$ with $k_\ell = k_{\ell+1}$ for $\ell \notin \text{supp} X$, one defines $\{[b_i, \ell_i]\}_{i=1}^{s(\mathbf{k})}$ as follows: For v such that $k_{1,v} \neq 0$ and for i such that $\sum_{j=0}^{v-1} k_{1,j} < i \leq \sum_{j=0}^v k_{1,j}$ let $b_i := v$. Let $m_1 > \dots > m_{k_{1,v}} > 0$ be the sequence of positive integers such that $k_{m_1,v} > k_{m_2,v} \dots > k_{m_{k_{1,v}},v}$ and such that $k_{m_j} > k_{m_j+1}$. Note that $m_j \in S$. Define $\ell_i := m_i - \sum_{j=0}^{v-1} k_{j,v}$. The detailed verification of the bijectivity of the map is tedious and we omit this. Instead we shall visualize the bijection below for Example 4. $\square$

[illegible]

TABLE 3. The 2-adic box and the minimal length r -tiling sequence

3. SUPERSINGULARITY CRITERION

Define the set of *supports* of X by $\text{supp}X := \{\ell \in \mathbb{Z} \mid c_\ell \neq 0\}$. For $r \geq 1$ let

$$\tilde{C}(r, \text{supp} X) := \sum_{M \in \tilde{K}(r, \text{supp} X)} \prod_{[b, \ell] \in M} c_\ell^{2^b} \in F.$$

Lemma 6. *Let notations be as above. Let λ be a rational number with $0 \leq \lambda \leq 1/2$.*

a) Suppose that for all $m \geq 1$, $n \geq 1$, and $1 \leq j \leq g$ one has

$$\tilde{s}(m2^{n+g-1} - j, \text{supp}X) \geq \lceil n\lambda \rceil.$$

Then

$$\text{NP}_1(X) \geq \lambda.$$

b) Suppose $\text{NP}_1(X) \geq \lambda$. Suppose there are positive integers $1 \leq j \leq g$, $n_0 \geq 1$ and $1 \leq m_0 \leq g/2$ such that

1) for all $m \geq 1$, $1 \leq n < n_0$ and for all $m > g/2$, $n = n_0$ we have

$$\tilde{s}(m2^{n+g-1} - j, \text{supp}X) \geq \lceil n\lambda \rceil;$$

2) $\tilde{s}(m_02^{n_0+g-1} - j, \text{supp}X) = \lceil n_0\lambda \rceil - 1$.

Then

$$\tilde{C}(m_02^{n_0+g-1} - j, \text{supp}X) = 0.$$

Proof. Let $W(F)$ be the ring of Witt vectors of F . Let $a_\ell \in W(F)$ be a lift of $c_\ell \in F$. Let $B := \min\{s(\mathbf{k}) \mid \mathbf{k} \in \mathbf{K}_r, \text{ and } k_\ell = k_{\ell+1} \text{ for all } \ell \notin \text{supp}X\}$. Set $0^0 := 1$. We recall $C_r(N)$ from [7]; we know from there that

$$C_r(N) \equiv \sum_{\mathbf{k} \in \mathbf{K}_r} 2^{s(\mathbf{k})} \prod_{\ell=1}^{d-1} a_\ell^{k_\ell - k_{\ell+1}} \pmod{2^{B+1}}.$$

Since $\text{ord}_2(C_r(N)) \geq B$, we may write

$$\frac{C_r(N)}{2^B} \equiv \sum_{\mathbf{k} \in \mathbf{K}_r} 2^{s(\mathbf{k})-B} \prod_{\ell=1}^{d-1} c_\ell^{k_\ell - k_{\ell+1}} \pmod{2}.$$

By the bijection discussed in section 2 we get

$$(2) \quad \left(\frac{C_r(N)}{2^{\tilde{s}(r, \text{supp}X)}} \pmod{2} \right) = \tilde{C}(r, \text{supp}X).$$

By (2), we note that a) and b) follow from the Key Lemma of [8]. (see also [7, Key-Lemma 2.1,i])). $\square$

Below we have two corollaries.

Corollary 7. Let X be as in (1). Let $\sigma := \max_{\ell \in \text{supp}X} s(\ell)$. Then we have $\text{NP}_1(X) \geq 1/\sigma \geq 1/d$.

Proof. For $r = m2^{n+g-1} - j$ consider any r -tiling sequence given by $m2^{n+g-1} - j = \sum_{i=1}^\nu \ell_i 2^{b_i}$. Clearly we have

$$\nu\sigma \geq \sum_{i=1}^\nu s(\ell_i) = \sum_{i=1}^\nu s(\ell_i 2^{b_i}) \geq s(m2^{n+g-1} - j) \geq n,$$

so $\nu \geq \lceil \frac{n}{\sigma} \rceil$. In particular

$$\tilde{s}(m2^{n+g-1} - j, \text{supp}X) \geq \lceil \frac{n}{\sigma} \rceil.$$

By Lemma 6 a) we have $\lambda \geq 1/\sigma$. But $c_d = 1 \neq 0$ so $\sigma \leq d$ and we have $\text{NP}_1(X) \geq 1/\sigma \geq 1/d$. $\square$

Remark 8. Any curve X given by an equation of the form $y^2 - y = \sum_{i=0}^s c_i x^{2^i+1}$ over F is supersingular. Indeed, apply the above corollary to this curve we get $\sigma = 2$ and so $\text{NP}_1(X) \geq 1/2$. (This result was proved in [10] in the case $F \subseteq \overline{\mathbb{F}_2}$.)

Corollary 9. *Suppose X given by equation (1) is supersingular. Let m_0, n_0 and j be positive integers satisfying hypothesis of Lemma 6 b) for $\lambda = \frac{1}{2}$. Suppose the set $\tilde{K}(m_0 2^{n_0+g-1} - j, \text{supp} X)$ consists of one element M . Then we have $\tilde{S}(M) \neq \{2g+1\}$ and if $\tilde{S}(M) = \{\ell, 2g+1\}$ with $1 \leq \ell \leq 2g$, then $c_\ell = 0$.*

Proof. Suppose $\tilde{S}(M) = \{2g+1\}$, then by Lemma 6 b) we have $\tilde{C}(m_0 2^{n_0+g-1} - j, \text{supp} X) = 0$. But it is easily seen that

$$\tilde{C}(m_0 2^{n_0+g-1} - j, \text{supp} X) = \prod_{[b, 2g+1] \in M} c_{2g+1}^{2^b} = 1$$

because $c_{2g+1} = 1$. Contradiction. This proves the first assertion.

Now suppose $\tilde{S}(M) = \{\ell, 2g+1\}$. By Lemma 6 again, we have

$$\begin{aligned} \tilde{C}(m_0 2^{n_0+g-1} - j, \text{supp} X) &= \prod_{[b, \ell] \in M} c_\ell^{2^b} \prod_{[b', 2g+1] \in M} c_{2g+1}^{2^{b'}} \\ &= \prod_{[b, \ell] \in M} c_\ell^{2^b} = c_\ell^{\sum_{[b, \ell] \in M} 2^b} = 0. \end{aligned}$$

Thus $c_\ell = 0$. □

4. SUPERSINGULARITY

We shall show that all curves in Table 1 are supersingular.

The supersingularity of X_8 follows from Remark 8. The supersingularity of the curve $C : y^2 - y = x^{33} + c_3 x^9 + c_1 x^3$ over F also follows from Remark 8. By observing that C covers X_5 , the supersingularity of X_5 follows.

Now we claim that X_6 is supersingular by a method suggested to us by Noam Elkies. We present it below: Let E be the elliptic curve defined by $y^2 - y = x^3 + t^{13} + c_3 t^3 + c_1 t$ over $F(t)$, which can be viewed as a quadratic twist of $E_0 : y^2 - y = x^3$ over the function field $F(X_6)$ of X_6 . The curve X_6 is supersingular if and only if the rank of $\text{Hom}_F(\text{Jac}(X_6), E_0)$, or equivalently the rank of $E(F(t))$ is 24, the maximal possible (see [9]). In the same vein as in [3, pages 3–4], one can show that the canonical height of a nonzero point in $E(F(t))$ is ≥ 8 . We shall show in the next paragraph that there are 196560 nonzero points with the minimal canonical height 8. Since there are no lattices of rank ≤ 23 with that many vectors of minimal length [1, page 23], this implies that $E(F(t))$ has rank 24 and the supersingularity of X_6 follows.

If $P = (x, y)$ is a point in $E(F(t))$ with x -coordinate of the form

$$x = a^{-16} t^6 + a^{-28} t^4 + a^{-40} t^2 + a^{32} t + x_0 + a^{-64} (t - t_0)^{-2},$$

for $a \in F^*$ and $x_0, t_0 \in F$, then P has canonical height 8 (see [2, Proposition 2]). Following closely the computations presented in [3, page 6–7], one finds that P lies in $E(F(t))$ if a, x_0 and t_0 satisfy the following equations:

$$\begin{aligned} (3) \quad & a^{4096} + c_3^{32} a^{1024} + c_1^{16} a^{256} + c_3^{16} a^{64} + c_1^4 a^{16} + c_3^2 a^4 + a = 0; \\ (4) \quad & a^{36} t_0^6 + a^{24} t_0^4 + a^{12} t_0^2 + a^{84} t_0 + a^{1092} + c_3^8 a^{324} + \\ & a^{312} + a^{156} + c_1^4 a^{132} + c_3^2 a^{120} + a^{117} + c_3 a^{84} + 1 = 0; \\ (5) \quad & a^{576} + a^{264} + c_3^4 a^{192} + (x_0^4 + x_0) a^{160} + a^{108} + t_0^8 = 0. \end{aligned}$$

(One may also check alternatively that the solutions to these 3 equations indeed yield x -coordinates of points in $E(F(t))$.) One can show that these 3 equations are

separable in a, t_0 and x_0 respectively by observing that their derivatives are nonzero constant. The number of nonzero solutions a of (3) is 4095. For each such a there are 6 solutions of t_0 satisfying (4). For each pair of (a, t_0) there are 4 solutions of x_0 satisfying (5). So there are totally $4095 \cdot 6 \cdot 4$ possible x -coordinates, each of which yields two points. This results in $2 \cdot 4 \cdot 6 \cdot 4095 = 196560$ points in $E(F(t))$.

5. NECESSARY

Lemma 6 b) shall find its extensive application here. A few remarks are in order. If there are indeed positive integers j, n_0, m_0 satisfying the hypothesis of Lemma 6 b) for some λ , then it is a finite problem to find them and verify. To see this, define $\tilde{T}(\nu, j, \text{supp}X)$ as the set of all sequences $\{[b_i, \ell_i]\}_{i=1}^\nu$ satisfying b) and that $\sum_{i=1}^\nu \ell_i 2^{b_i} \equiv -j \pmod{2^{b_\nu}}$. Clearly, $\tilde{T}(\nu, j, \text{supp}X)$ is a finite set, and if $\nu = \tilde{s}(m_0 2^{n_0+g-1} - j, \text{supp}X)$ for some m_0 and n_0 , then $\tilde{T}(\nu, j, \text{supp}X)$ contains $\tilde{K}(m_0 2^{n_0+g-1} - j, \text{supp}X)$ for every m with

$$\tilde{s}(m_0 2^{n_0+g-1} - j, \text{supp}X) \leq \tilde{s}(m_0 2^{n_0+g-1} - j, \text{supp}X).$$

So one can verify if b) is satisfied by computing $\tilde{T}(\nu, j, \text{supp}X)$ for $1 \leq j \leq g$ and for $\nu \geq 1$, and check for each such ν and j whether there exist m_0 and n_0 with the required property.

We shall verify the hypotheses of b) by the aforementioned algorithm, together with a few isolated tricks to accelerate the program. For instance, we use an observation that if two sequences $\{[b_i, \ell_i]\}_{i=1}^\nu$ and $\{[b'_i, \ell'_i]\}_{i=1}^{\nu'}$ satisfy $\sum_{i=1}^\nu \ell_i 2^{b_i} = \sum_{i=1}^{\nu'} \ell'_i 2^{b'_i}$ and $\nu' < \nu$, then one can ignore all sequences including $\{[b_i, \ell_i]\}_{i=1}^\nu$ as a subsequence.

For convenience of the reader we first quote Proposition 4.1 and Theorem 1.1 of [7] below.

Proposition 10. *Let X be a hyperelliptic curve over F of genus g and 2-rank zero. Let $m < 2g$ be an odd integer, such that the binomial coefficient $\binom{2g+1}{2^k m}$ is odd for some $k \geq 0$. Then there exists an equation for X of the form*

$$(6) \quad y^2 - y = \sum_{i=0}^g c_{2i+1} x^{2i+1}$$

with $c_{2g+1} = 1$ and $c_m = 0$. Moreover, there are only finitely many equations for X of this form.

Theorem 11. *Let $g \geq 3$. Write $h = \lfloor \log_2(g+1) + 1 \rfloor$. Let X be a hyperelliptic curve over F given by an equation $y^2 - y = \sum_{i=1}^{2g+1} c_i x^i$ where $c_{2g+1} = 1$. Then*

- (I) $\text{NP}_1(X) \geq \frac{1}{h}$.
- (II) If $g < 2^h - 2$ and $c_{2^h-1} \neq 0$ then $\text{NP}_1(X) = \frac{1}{h}$.
- (III) If $g = 2^h - 2$ and $c_{2^h-1} \neq 0$ or $c_{3 \cdot 2^{h-1}-1} \neq 0$ then $\text{NP}_1(X) = \frac{1}{h}$.

Now we will proceed with the proof of Theorem 1 for genus $g \geq 5$. For the rest of the proof we shall suppress the $\text{supp}X$ from the notations $\tilde{C}(r, \text{supp}X)$ and $\tilde{K}(r, \text{supp}X)$. For every genus we start with a curve with an equation in a box. Then we use supersingularity criterion to derive vanishness of coefficients of this initial equation, which accumulate to our desired equation declared in Table 1.

5.1. $g = 5$. Suppose X is a hyperelliptic supersingular curve over F of genus 5. By Theorem 11 II) and Proposition 10 it has an equation

$$y^2 - y = x^{11} + c_9 x^9 + c_3 x^3 + c_1 x$$

for some $c_1, c_3, c_9 \in F$.

Let $\text{supp}X = \{1, 3, 9, 11\}$. Let $g = 5$. We checked that for $n_0 = 13$, $m_0 = 1$ and $j = 4$ the conditions of Lemma 6 b) are satisfied, and $\tilde{K}(m_0 2^{n_0+g-1} - j)$ consists of a unique element M as follows

$$2^{17} - 4 = 11 \cdot 2^{13} + 9 \cdot 2^{12} + 11 \cdot 2^8 + 9 \cdot 2^7 + 11 \cdot 2^3 + 9 \cdot 2^2.$$

Note that $\tilde{S}(M) = \{9, 11\}$. (See Example 4.) Then Corollary 9 implies that $c_9 = 0$.

5.2. $g = 6$. Suppose X is a hyperelliptic supersingular curve over F of genus 6. By Theorem 11 III) and Proposition 10 it has an equation

$$y^2 - y = x^{13} + c_9 x^9 + c_3 x^3 + c_1 x$$

for some $c_1, c_3, c_9 \in F$.

Let $\text{supp}X = \{1, 3, 9, 13\}$. Let $g = 6$. We checked that for $n_0 = 17$, $m_0 = 1$, $j = 4$ the conditions of Lemma 6 b) are satisfied, and $\tilde{K}(m_0 2^{n_0+g-1} - 4)$ consists of a unique element M as follows

$$2^{22} - 4 = 9 \cdot 2^{18} + 13 \cdot 2^{17} + 9 \cdot 2^{13} + 13 \cdot 2^{12} + 9 \cdot 2^8 + 13 \cdot 2^7 + 9 \cdot 2^3 + 13 \cdot 2^2.$$

Note that $\tilde{S}(M) = \{9, 13\}$. By Corollary 9 we have $c_9 = 0$.

5.3. $g = 8$. Suppose X is a hyperelliptic supersingular curve over F of genus 8. By Theorem 11 II) and Proposition 10 it has an equation

$$y^2 - y = x^{17} + c_{13} x^{13} + c_{11} x^{11} + c_9 x^9 + c_7 x^7 + c_5 x^5 + c_3 x^3$$

for some $c_\ell \in F$.

Let $\text{supp}X = \{3, 5, 7, 9, 11, 13, 17\}$. We found for $k = 0, 1, 2, 3, 4$ and 5 that $\tilde{K}(2^{21-3k} - 8)$ consists of sequences $\{[b_i, \ell_i]\}_{i=1}^{6-k}$ with either $[b_{6-k}, \ell_{6-k}] = [18-3k, 7]$ or $[b_{6-k}, \ell_{6-k}] = [17-3k, 13]$ and $[b_{5-k}, \ell_{5-k}] = [15-3k, 11]$. Hence we get

$$\begin{aligned} \tilde{C}(2^{21} - 8) &= c_7^{2^{18}} \tilde{C}(2^{18} - 8) + (c_{11} c_{13}^4)^{2^{15}} \tilde{C}(2^{15} - 8) \\ \tilde{C}(2^{18} - 8) &= c_7^{2^{15}} \tilde{C}(2^{15} - 8) + (c_{11} c_{13}^4)^{2^{12}} \tilde{C}(2^{12} - 8) \\ \tilde{C}(2^{15} - 8) &= c_7^{2^{12}} \tilde{C}(2^{12} - 8) + (c_{11} c_{13}^4)^{2^{9}} \tilde{C}(2^9 - 8) \\ \tilde{C}(2^{12} - 8) &= c_7^{2^9} \tilde{C}(2^9 - 8) + (c_{11} c_{13}^4)^{2^{6}} \tilde{C}(2^6 - 8) \\ \tilde{C}(2^9 - 8) &= c_7^{2^6} \tilde{C}(2^6 - 8) + (c_{11} c_{13}^4)^{2^{3}} \tilde{C}(2^3 - 8) \\ \tilde{C}(2^6 - 8) &= c_7^{2^3} \tilde{C}(2^3 - 8). \end{aligned}$$

We checked that the conditions of Lemma 6 b) are satisfied for $\lambda = \frac{1}{2}$, $n_0 = 11$, $m_0 = 1$ and $j = 8$. Hence $\tilde{C}(2^{18} - 8) = 0$. We also checked that they are satisfied for $\lambda = \frac{4}{9}$, $n_0 = 14$, $m_0 = 1$ and $j = 8$. Hence $\tilde{C}(2^{21} - 8) = 0$. All these equations imply $c_7 = 0$ and $c_{11} c_{13} = 0$.

Suppose $c_{11} = 0$. Let $\text{supp}X = \{3, 5, 9, 13, 17\}$. We checked that for $\lambda = \frac{1}{2}$, $j = 4$, $n_0 = 19$ and $m_0 = 1$ the conditions of Lemma 6 b) are satisfied, and $\tilde{K}(m_0 2^{n_0+g-1} - 4)$ consists of a unique element M as follows

$$\begin{aligned} 2^{26} - 4 &= 13 \cdot 2^{22} + 17 \cdot 2^{19} + 13 \cdot 2^{18} + 13 \cdot 2^{14} + 17 \cdot 2^{11} + 13 \cdot 2^{10} \\ &\quad + 13 \cdot 2^6 + 17 \cdot 2^3 + 13 \cdot 2^2. \end{aligned}$$

Note that $\tilde{S}(M) = \{13, 17\}$. By Corollary 9 we have $\boxed{c_{13} = 0}$.

Suppose $c_{13} = 0$. Let $\text{supp}X = \{3, 5, 9, 11, 17\}$. We checked that for $\lambda = \frac{1}{2}$, $j = 4$, $n_0 = 19$ and $m_0 = 1$ the conditions of Lemma 6 b) are satisfied, and $\tilde{K}(m_0 2^{n_0+g-1} - 4)$ consists of a unique element M as follows

$$\begin{aligned} 2^{26} - 4 &= 11 \cdot 2^{22} + 17 \cdot 2^{20} + 11 \cdot 2^{18} + 11 \cdot 2^{14} + 17 \cdot 2^{12} + 11 \cdot 2^{10} + \\ &\quad 11 \cdot 2^6 + 17 \cdot 2^4 + 11 \cdot 2^2. \end{aligned}$$

Note that $\tilde{S}(M) = \{11, 17\}$. By Corollary 9 we have $\boxed{c_{11} = 0}$.

5.4. $g = 9$. Suppose X is a hyperelliptic supersingular curve over F of genus 9. By Theorem 11 II and Proposition 10 it has an equation

$$\boxed{y^2 - y = x^{19} + c_{13}x^{13} + c_{11}x^{11} + c_9x^9 + c_7x^7 + c_5x^5 + c_3x^3 + c_1x}$$

for some $c_\ell \in F$.

Let $\text{supp}X = \{1, 3, 5, 7, 9, 11, 13, 19\}$. We found for $k = 0, \dots, 7$ that $\tilde{K}(2^{27-3k} - 8)$ consists of all sequences $\{[b_i, \ell_i]\}_{i=1}^{8-k}$ possessing the following property:

$$\left\{ \begin{array}{l} [b_{8-k}, \ell_{8-k}] = [24 - 3k, 7] \text{ or} \\ [b_{8-k}, \ell_{8-k}] = [23 - 3k, 13] \text{ and } [b_{7-k}, \ell_{7-k}] = [21 - 3k, 11] \text{ or} \\ [b_{8-k}, \ell_{8-k}] = [23 - 3k, 11] \text{ and } [b_{7-k}, \ell_{7-k}] = [21 - 3k, 19] \text{ or} \\ [b_{8-k}, \ell_{8-k}] = [23 - 3k, 13] \text{ and } [b_{7-k}, \ell_{7-k}] = [20 - 3k, 19] \text{ and } [b_{6-k}, \ell_{6-k}] = [18 - 3k, 19]. \end{array} \right.$$

Hence

$$\begin{aligned} \tilde{C}(2^{27-3k} - 8) &= c_7^{2^{24-3k}} \tilde{C}(2^{24-3k} - 8) + (c_{11}c_{13}^4 + c_{11}^4)^{2^{21-3k}} \tilde{C}(2^{21-3k} - 8) \\ &\quad + c_{13}^{2^{23-3k}} \tilde{C}(2^{18-3k} - 8). \end{aligned}$$

We checked that the conditions of Lemma 6 b) are satisfied for $\lambda = \frac{1}{2}$, $j = 8$ and (n_0, m_0) equal to $(17, 4)$, $(15, 2)$ and $(13, 1)$. Using the same kind of reasoning as for genus 8, one concludes that $\boxed{c_7 = 0}$, $\boxed{c_{13} = 0}$ and $c_{11}c_{13}^4 + c_{11}^4 = 0$, which clearly implies $\boxed{c_{11} = 0}$.

So we continue with $X = \{1, 3, 5, 9, 19\}$. We checked that for $n_0 = 19$, $m_0 = 1$, $j = 8$ the conditions of Lemma 6 b) are satisfied, and $\tilde{K}(m_0 2^{n_0+g-1} - 8)$ consists of a unique element M as follows

$$2^{27} - 8 = 5 \cdot 2^{24} + 19 \cdot 2^{21} + 19 \cdot 2^{19} + 5 \cdot 2^{16} + 19 \cdot 2^{13} + 19 \cdot 2^{11} + 5 \cdot 2^8 + 19 \cdot 2^5 + 19 \cdot 2^3.$$

Note that $\tilde{S}(M) = \{5, 19\}$. By Corollary 9 we have $\boxed{c_5 = 0}$.

Now set $\text{supp}X = \{1, 3, 9, 19\}$. We checked that for $n_0 = 25$, $m_0 = 3$, $j = 8$ the conditions of Lemma 6 b) are satisfied, and $\tilde{K}(m_0 2^{n_0+g-1} - 8)$ consists of a unique element M as follows

$$\begin{aligned} 3 \cdot 2^{33} - 8 &= 19 \cdot 2^{30} + 19 \cdot 2^{28} + 3 \cdot 2^{25} + 19 \cdot 2^{23} + 3 \cdot 2^{20} + 19 \cdot 2^{18} + 3 \cdot 2^{15} \\ &\quad + 19 \cdot 2^{13} + 3 \cdot 2^{10} + 19 \cdot 2^8 + 3 \cdot 2^5 + 19 \cdot 2^3. \end{aligned}$$

Note that $\tilde{S}(M) = \{3, 19\}$. By Corollary 9 we have $\boxed{c_3 = 0}$.

Finally, set $\text{supp}X = \{1, 9, 19\}$. We found that for $n_0 = 35$, $m_0 = 3$, $j = 8$ and $\lambda = \frac{1}{2}$ the conditions of Lemma 6 b) are satisfied, and that $\tilde{K}(3 \cdot 2^{43} - 8)$ consists of sequences $\{[b_i, \ell_i]\}_{i=1}^{17}$ with $[b_{17}, \ell_{17}] = [40, 19]$, $[b_{16}, \ell_{16}] = [38, 19]$ and for $k = 1, \dots, 5$ we either have $[b_{3k}, \ell_{3k}] = [7k - 4, 19]$, $[b_{3k+1}, \ell_{3k+1}] = [7k - 2, 19]$, $[b_{3k+2}, \ell_{3k+2}] = [7k + 1, 1]$ or $[b_{3k}, \ell_{3k}] = [7k - 4, 19]$, $[b_{3k+1}, \ell_{3k+1}] = [7k - 2, 9]$, $[b_{3k+2}, \ell_{3k+2}] = [7k - 1, 9]$. So from Lemma 6 it follows that

$$0 = \tilde{C}(3 \cdot 2^{43-8} - 8) = (c_1^{32} + c_9^{12})^{2^3+2^{10}+2^{17}+2^{24}+2^{31}}.$$

Thus $\boxed{c_9^3 = c_1^8}$, and we conclude that every hyperelliptic supersingular curve of genus 9 over F has to have an equation of the form $y^2 - y = x^{19} + c^8 x^9 + c^3 x$ for some $c \in F$.

A straightforward computation shows that the curve X with equation $y^2 - y = x^{19} + c^8 x^9 + c^3 x$ over $\overline{\mathbb{F}}_2$ is supersingular for $c \in \mathbb{F}_2$ but is *not* supersingular for $c \in \mathbb{F}_{2^2} - \mathbb{F}_2$. Therefore, this curve is supersingular for only finitely many $c \in \overline{\mathbb{F}}_2$. In other words, the open locus of hyperelliptic supersingular curves of genus 9 over $\overline{\mathbb{F}}_2$ is of dimension 0.

Let $c \in F - \overline{\mathbb{F}}_2$. Suppose $X : y^2 - y = x^{19} + c^8 x^9 + c^3 x$ is supersingular. Then its specialization yields infinitely many supersingular curves over $\overline{\mathbb{F}}_2$ because the Grothendieck Specialization theorem says that the Newton polygon goes up under specialization maps (see [5]). This would contradict the conclusion in the previous paragraph. Therefore, X has to be defined over $\overline{\mathbb{F}}_2$.

Question 12. What are normal forms of these hyperelliptic supersingular curve of genus 9 over $\overline{\mathbb{F}}_2$? Are $y^2 - y = x^{19}$ and $y^2 - y = x^{19} + x^9 + x$ the only ones?

Computations by our algorithm reveal some pattern of the dimension of the open locus $\mathcal{HS}_g/\overline{\mathbb{F}}_2$ of hyperelliptic supersingular curves of genus $g \geq 10$ over $\overline{\mathbb{F}}_2$. Vaguely speaking $\dim(\mathcal{HS}_g/\overline{\mathbb{F}}_2)$ is related to $s(2g + 1)$ in the sense that $\dim(\mathcal{HS}_g/\overline{\mathbb{F}}_2)$ goes smaller as the ratio of 1's in the binary expansion of $2g + 1$ goes bigger. This is already evident in our previous paper [7] which shows that for $2g + 1 = 2^h - 1$ for some $h > 2$ then $\mathcal{HS}_g/\overline{\mathbb{F}}_2$ is empty. This paper further explores this relationship.

Question 13. It is desirable to formulate the pattern of $\dim(\mathcal{HS}_g/\overline{\mathbb{F}}_2)$ more precisely by more numerical data.

REFERENCES

- [1] J. CONWAY; N. SLOANE: *Sphere packings, lattices and groups*, 3rd ed., Springer-Verlag, New York, 1993.
- [2] NOAM D. ELKIES: Mordell-Weil lattices in characteristic 2: I. Construction and first properties, *International Mathematics Research Notices* no. **8** (1994), 343–361.
- [3] NOAM D. ELKIES: Mordell-Weil lattices in characteristic 2: II. The Leech lattice as a Mordell-Weil lattice. *Invent. Math.* **128** (1997), 1–8.
- [4] J. DE JONG; FRANS OORT: Hyperelliptic curves in abelian varieties. *Algebraic geometry*, 5. J. Math. Sci. **82** (1996), 3211–3219.
- [5] NICHOLAS KATZ: Slope filtration of F-crystals, *Astérisque* **63** (1979), 113–164.
- [6] KE-ZHENG LI; FRANS OORT: *Moduli of supersingular abelian varieties*, Lecture notes in mathematics, **1680**. Springer. 1998.
- [7] JASPER SCHOLTEN; HUI JUNE ZHU: Hyperelliptic curves in characteristic 2, electronic preprint available online at <http://xxx.lanl.gov/abs/math.AG/0012178>, 2001.
- [8] JASPER SCHOLTEN; HUI JUNE ZHU: Slope estimates of Artin-Schreier curves, electronic preprint available online at <http://xxx.lanl.gov/abs/math.AG/0105005>, 2001.
- [9] I. SHAFAREVICH; J. TATE: The rank of elliptic curves, *Soviet Math. Dokl.* **8** (1967), 917–920.

- [10] GERARD VAN DER GEER; MARCEL VAN DER VLUGT: On the existence of supersingular curves of given genus. *J. Reine Angew. Math.*, **458** (1995), 53–61.
- [11] GERARD VAN DER GEER; MARCEL VAN DER VLUGT:
- [12] HUI JUNE ZHU: p -adic variation of L functions of exponential sums, I electronic preprint available online at <http://xxx.lanl.gov/abs/math.AG/0111194>, 2001.

MATHEMATISCH INSTITUUT, KATHOLIEKE UNIVERSITEIT NIJMEGEN, POSTBUS 9010, 6500 GL NIJMEGEN. THE NETHERLANDS.

E-mail address: `scholten@sci.kun.nl`

DEPARTMENT OF MATHEMATICS, UNIVERSITY OF CALIFORNIA, BERKELEY, CA 94720-3840. THE UNITED STATES.

E-mail address: `zhu@alum.calberkeley.org`